


Cembranoids from Soft Coral *Sarcophyton tortuosum*

Kuan-Hua Lin (林官嬌), Jyh-Horng Sheu (許志宏)^{*}

Department of Marine Resources, National Sun Yat-Sen University, Kaoshiung, 804, Taiwan, ROC

Keywords: marine, soft coral, Cembranoids, diterpenes

Four new cembranoids, collected at Orchid island, were isolated from the soft coral *Sarcophyton tortuosum*, together with four known cembranoids. The structures and relative stereochemistry of the compounds were determined by detailed spectroscopic analysis including 1D and 2D NMR (¹H, ¹³C, ¹H-¹H COSY, HSQC, HMBC, and NOESY) data in association with MS experiments.



Reference:

1. Zhang, C.; Li, J.; Su, J.; Liang, Y.; Yang, X.; Zheng, K.; Zeng, L. *J. Nat. Prod.* **2006**, *69*, 1476–1480.
2. Uchio, Y.; Nitya, M.; Twaoawa, T. *Chem. Lett.* **1983**, 613–616.
3. Toth, J. A.; Burreson, B. J.; Clardy, J.; Finermoore, J.; Scheuer, P. J. *Tetrahedron*, **1980**, *36*, 1307-1309.
4. 林好芳, 野外型與養殖型柔軟指形軟珊瑚所含天然物之研究, 國立中山大學海洋生物科技暨資源學系博士論文, 中華民國九十七年。